arXiv:1103.4058v1 [math.AP] 21 Mar 2011

NONSTATIONARY FLOW FOR THE NAVIER-STOKES EQUATIONS IN A CYLINDRICAL PIPE

JOANNA RENCŁAWOWICZ & WOJCIECH M. ZAJĄCZKOWSKI

Abstract. In cylindrical domain, we consider the nonstationary flow with prescribed inflow and outflow, modelled with Navier-Stokes equations under the slip boundary conditions. Using smallness of some derivatives of inflow function, external force and initial velocity of the flow, but with no smallness restrictions on the inflow, initial velocity neither force, we prove existence of solutions in $W_2^{2,1}$.

1. Introduction

In the paper we study viscous incompressible fluid motion in a finite cylinder with large inflow and outflow. The nonstationary flow is described by Navier-Stokes equations and we impose the boundary slip conditions, with friction between the flux and the part of the boundary. Namely, we consider the following initial boundary value problem.

$$\begin{aligned}
&v_t + v\cdot\nabla v - \operatorname{div}\mathbb{T}(v,p) = f && \text{in } \ \Omega^T = \Omega\times(0,T),\\
&\operatorname{div} v = 0 && \text{in } \ \Omega^T,\\
&v\cdot\bar n = 0 && \text{on } S_1^T,\\
&\nu\bar n\cdot\mathbb{D}(v)\cdot\bar\tau_\alpha + \gamma v\cdot\bar\tau_\alpha = 0, \quad \alpha=1,2, && \text{on } \ S_1^T,\\
&v\cdot\bar n = d && \text{on } S_2^T,\\
&\bar n\cdot\mathbb{D}(v)\cdot\bar\tau_\alpha = 0, \quad \alpha=1,2, && \text{on } S_2^T,\\
&v\big|_{t=0} = v(0) && \text{in } \ \Omega,
\end{aligned}\tag{1.1}$$

where $\Omega\subset\mathbb{R}^3$ is a cylindrical domain, $S=\partial\Omega$, v is the velocity of the fluid motion with $v(x,t)=(v_1(x,t),v_2(x,t),v_3(x,t))\in\mathbb{R}^3$, $p=p(x,t)\in\mathbb{R}^1$ denotes the pressure, $f=f(x,t)=(f_1(x,t),f_2(x,t),f_3(x,t))\in\mathbb{R}^3$ – the external force field, $x=(x_1,x_2,x_3)$ are the Cartesian coordinates, $\bar n$ is the unit outward vector normal to the boundary S and $\bar\tau_\alpha$, $\alpha=1,2$, are tangent vectors to S and $\cdot$ denotes the scalar product in $\mathbb{R}^3$. $\mathbb{T}(v,p)$ is the stress tensor of the form

$$\mathbb{T}(v,p)=\nu\mathbb{D}(v)-p\mathbb{I},$$

Date: January 15, 2013.

2010 *Mathematics Subject Classification.* Primary 35Q30; Secondary 76D03, 76D05.

Key words and phrases. Navier-Stokes equation, Neumann boundary-value problem, Dirichlet boundary-value problem, regular solutions, large flux, slip boundary conditions.

Research supported by MNiSW grant no N N201 396937.

where ν is the constant viscosity coefficient and $\mathbb{I}$ is the unit matrix. Next, $\gamma > 0$ is the slip coefficient and $\mathbb{D}(v)$ denotes the dilatation tensor of the form

$$\mathbb{D}(v) = \{v_{i,x_j} + v_{j,x_i}\}_{i,j=1,2,3}.$$

We consider $\Omega \subset \mathbb{R}^3$ which is a cylindrical type domain parallel to the axis x_3 with arbitrary cross section. We split S into S_1 and S_2 so that $S = S_1 \cup S_2$ where S_1 is the part of the boundary which is parallel to the axis x_3 and S_2 is perpendicular to x_3. Consequently,

$$\begin{aligned} S_1 &= \{x \in \mathbb{R}^3 : \varphi_0(x_1, x_2) = c_0,\ -a < x_3 < a\}, \\ S_2(-a) &= \{x \in \mathbb{R}^3 : \varphi_0(x_1, x_2) < c_0,\ \ x_3 = -a\}, \\ S_2(a) &= \{x \in \mathbb{R}^3 : \varphi_0(x_1, x_2) < c_0,\ \ x_3 = a\} \end{aligned}$$

where a, c_0 are positive given numbers and $\varphi_0(x_1, x_2) = c_0$ describes a sufficiently smooth closed curve in the plane $x_3 = \text{const}$.



Figure 1. Domain Ω.

To describe inflow and outflow we define

$$\begin{aligned} d_1 &= -v \cdot \bar{n}|_{S_2(-a)} \\ d_2 &= v \cdot \bar{n}|_{S_2(a)} \end{aligned} \tag{1.2}$$

with $d_i \geq 0, i = 1, 2$. Using $(1.1)_{2,3}$ and (1.2) we conclude the following compatibility condition

$$\Phi \equiv \int_{S_2(-a)} d_1 dS_2 = \int_{S_2(a)} d_2 dS_2, \tag{1.3}$$

where Φ is the flux.

The main goal of the paper is to show the long time existence of regular solutions to problem (1.1) with arbitrary large external force f, initial velocity $v(0)$ and inflow-outflow d. We can obtain the existence result by regularizing weak solutions. In general, we follow the ideas from [RZ1, Z1].

Regularity of solutions for Navier-Stokes equations, even with no flux, requires some smallness conditions. Many results of this type were proved assuming smallness of initial velocity, some restrictions on domain (so called thin domain, $\Omega = \Omega' \times (0, \varepsilon), \Omega' \in \mathbb{R}^2$ with small ε) or special structure of solutions (so that the solution is close to 2-dimensional solution). We mention here as well some results in case of stationary or nonstationary flux, with the same type of boundary condition that we assume: slip condition on the part of boundary, involving or neglecting friction between the boundary and the fluid. In [M1], the existence for large data is obtained under some geometrical constraints for 2d model in steady and evolutionary case. In [M2], steady Navier Stokes equations in pipe-like domain are investigated and existence is shown for a class of cylindrical symmetric solutions. In [Z4], the problem of nonstationary flow in axially symmetric domain is examined and the result concerns the existence for solutions close to axially symmetric solutions and the inflow and outflow sufficiently close to homogeneous flux.

We show the existence of regular solutions in a class of flows with arbitrary large initial velocity $v(0)$, inflow d_1 and external force f. Instead, we use smallness of derivatives of initial velocity and force in the direction along the x_3 axis and smallness of some derivatives of the inflow.

To formulate the main result, we introduce some notation.

We will use Sobolev spaces

$$W_p^{s,s/2}(Q^T), \quad Q \in \{\Omega, S\}, \quad s \in \mathbb{Z}_+ \cup \{0\}, \quad p \in [1,\infty],$$

with the following norm for even s

$$W_p^{s,s/2}(Q^T) = \{u : \|u\|_{W_p^{s,s/2}(Q^T)} = \left(\sum_{|\alpha|+2a\le s}\int_{Q^T}|D_x^\alpha\partial_t^a u|^p dxdt\right)^{1/p} \le \infty\},$$

where $D_x^\alpha = \partial_{x_1}^{\alpha_1}\partial_{x_2}^{\alpha_2}\partial_{x_3}^{\alpha_3}$, $\alpha = (\alpha_1,\alpha_2,\alpha_3)$, $|\alpha| = \alpha_1+\alpha_2+\alpha_3$, $\alpha_i \in \mathbb{Z}_+ \cup \{0\}$, $i = 1,2,3$. In particular,

$$H^s(Q) = W_2^s(Q).$$

We will need also

$$V_2^0(\Omega^T) = \{u : ||u||_{V_2^0(\Omega^T)} = \operatorname{ess}\sup_{t\in(0,T)}||u||_{L_2(\Omega)} + \left(\int_0^T ||\nabla u||^2_{L_2(\Omega)}dt\right)^{1/2} < \infty\},$$

$$V_2^1(\Omega^T) = \{u : ||u||_{V_2^1(\Omega^T)} = \operatorname{ess}\sup_{t\in(0,T)}||u||_{H^1(\Omega)} + \left(\int_0^T ||\nabla u||^2_{H^1(\Omega)}dt\right)^{1/2} < \infty\}.$$

Let A be a bound for the V_0^2 norm of v – a weak solution of (1.1), i.e.

$$\|v\|^2_{V_2^0(\Omega^t)} + \gamma\sum_{\alpha=1}^2\int_0^t \|v\cdot\bar\tau_\alpha\|^2_{L_2(S_1)} \le A^2$$

where the existence of such constant can be obtained as it is shown in [RZ1], see Lemma 2.1 below.

Definition 1.1. *We define*

$$D_0 = \|d\|^6_{L_\infty(0,t;L_3(S_2))} + A^2 + 1, \tag{1.4}$$

$$\begin{aligned} \Lambda_2 &= \|(\operatorname{rot} f)_3\|_{L_2(0,t;H^2(\Omega))} + \|(\operatorname{rot} v)_3(0)\|_{L_2(\Omega)} + 1 \\ \Lambda_3 &= \|f\|_{L_{5/3}(\Omega^T)} + \|v(0)\|_{W^{4/5}_{5/3}(\Omega)}, \\ \Lambda_4 &= \|f\|_{L_2(\Omega^T)} + \|v(0)\|_{H^1(\Omega)}. \end{aligned} \tag{1.5}$$

We note $h = v_{,x_3}$, $g = f_{,x_3}$, $x' = (x_1,x_2)$. *We set*

$$\Lambda = D_1 + D_2 + \|f_3\|^2_{L_2(0,t;L_{4/3}(S_2))} + \|g\|^2_{L_2(0,t;L_{6/5}(\Omega))} + \|h(0)\|^2_{L_2(\Omega)}, \tag{1.6}$$

where

$$
\begin{aligned}
&D_1 = \|d_t\|^2_{L_2(0,t;H^1(S_2))} + \|d_{,x'}\|^2_{L_2(0,t;H^1(S_2))},\\
&D_2 = \|d_{,x'}\|^2_{L_\infty(0,t;H^1(S_2))}.
\end{aligned}
\tag{1.7}
$$

Condition 1. *Let $D_0, \Lambda, \Lambda_2, \Lambda_3, \Lambda_4$ be finite.*

Condition 2. *Let Λ be so small and $\|g\|_{L_2(\Omega^T)}$, $\|h(0)\|_{H^1(\Omega)}$ be such that there exists constant $\mathcal{A}$ satisfying the relation*

$$
\begin{aligned}
&cD_0^2\Lambda^2\left[(\mathcal{A}+\Lambda)+c(\mathcal{A}+\mathcal{A}^2)+c\Lambda_2+c\Lambda_3\right]\\
&\exp\left[\int_0^t (\|d\|^6_{L_3(S_2)}dt + cD_0(\mathcal{A}+\Lambda)+c(\mathcal{A}+\mathcal{A}^2)+c\Lambda_2+c\Lambda_3\right]\\
&\qquad +\|g\|^2_{L_2(\Omega^T)} + \|h(0)\|^2_{H^1(\Omega)} \le \mathcal{A}^2
\end{aligned}
$$

Theorem 1. *Assume that Conditions 1 and 2 hold. Then there exists a solution to problem (1.1) such that*

$$
v, v_{,x_3} \in W_2^{2,1}(\Omega^T), \nabla p, \nabla p_{,x_3} \in L_2(\Omega^T)
$$

and

$$
\begin{aligned}
&\|v_{,x_3}\|_{W_2^{2,1}(\Omega^T} \le \mathcal{A},\\
&\|v\|_{W_2^{2,1}(\Omega^T} + \|\nabla p\|_{L_2(\Omega^T)} \le \varphi(\mathcal{A}, D_0, \Lambda, \Lambda_2, \Lambda_3, \Lambda_4),\\
&\|\nabla p_{,x_3}\|_{L_2(\Omega^T)} \le \varphi(\mathcal{A}, D_0, \Lambda, \Lambda_2, \Lambda_3, \Lambda_4),
\end{aligned}
\tag{1.8}
$$

where φ is an increasing positive function.

The structure of the paper is the following. First, in Section 2, since the boundary conditions in the problem (1.1) are not homogeneous, we introduce auxiliary variables and study the corresponding equations. In Section 3.1 and 3.2 we obtain the energy inequalities for $h = v_{,x_3}$ and next for $\chi = (\operatorname{rot} v)_3$. Then, through the elliptic system relating χ and v_{x_1}, v_{x_2}, we can conclude the relation for v in terms of h norms in Section 3.3. Next, the energy inequality for h is improved by getting a bound for the energy norm of h, depending on some norm of ∇v. This relation is used to show the estimate for h and v. In Section 4, we prove the existence using a priori estimates and the Leray-Schauder theorem.

2. Auxiliary problems

In order to define weak solutions to problem (1.1) and to obtain the energy type estimate we need to make the boundary condition $(1.1)_4$ homogeneous. Thus, we have to reformulate the main problem using some auxiliary functions.

First, we extend functions corresponding to inflow and outflow so that

$$
\tilde{d}_i|_{S_2(a_i)} = d_i,\ i = 1, 2,\ a_1 = -a,\ a_2 = a
\tag{2.1}
$$

We introduce the function η, see [L].

$$
\eta(\sigma; \varepsilon, \rho) = \begin{cases} 1 & 0 \le \sigma \le \rho e^{-1/\varepsilon} \equiv r,\\ -\varepsilon \ln \dfrac{\sigma}{\rho} & r < \sigma \le \rho,\\ 0 & \rho < \sigma < \infty. \end{cases}
$$

We calculate

$$\frac{d\eta}{d\sigma} = \eta'(\sigma;\varepsilon,\rho) = \begin{cases} 0 & 0<\sigma\le r,\\ -\dfrac{\varepsilon}{\sigma} & r<\sigma\le\rho,\\ 0 & \rho<\sigma<\infty. \end{cases}$$

so that $|\eta'(\sigma;\varepsilon,\rho)| \le \dfrac{\varepsilon}{\sigma}$. We define functions η_i on the neighborhood of S_2 (inside Ω):

$$\eta_i = \eta(\sigma_i;\varepsilon,\rho),\ \ i=1,2,$$

where σ_i denote local coordinates defined on small neighborhood of $S_2(a_i)$:

$$\sigma_1 = a + x_3,\ \ \sigma_2 = a - x_3$$

and we set

$$\begin{aligned} \alpha &= \sum_{i=1}^{2} \tilde{d}_i\eta_i,\\ b &= \alpha\bar{e}_3,\ \ \bar{e}_3 = (0,0,1). \end{aligned} \tag{2.2}$$

We construct function u so that

$$u = v - b. \tag{2.3}$$

Therefore,

$$\begin{aligned} \operatorname{div} u &= -\operatorname{div} b = -\alpha_{x_3} \quad \text{in} \ \ \Omega,\\ u\cdot\bar{n} &= 0 \quad \text{on} \ \ S. \end{aligned}$$

Then, the boundary condition for u is homogeneous. The compatibility condition takes the form

$$\int_\Omega \alpha_{,x_3}\,dx = -\int_{S_2(-a)} \alpha|_{x_3=-a} dS_2 + \int_{S_2(a)} \alpha|_{x_3=a} dS_2 = 0$$

We define function φ as a solution to the Neumann problem

$$\begin{aligned} \Delta\varphi &= -\operatorname{div} b \quad \text{in} \ \ \Omega,\\ \bar{n}\cdot\nabla\varphi &= 0 \quad \text{on} \ \ S,\\ \int_\Omega \varphi dx &= 0. \end{aligned} \tag{2.4}$$

Next, we set

$$w = u - \nabla\varphi = v - (b + \nabla\varphi) \equiv v - \delta. \tag{2.5}$$

Then, (w,p) is a solution to the following problem

$$
\begin{aligned}
&w_t + w\cdot\nabla w + w\cdot\nabla\delta + \delta\cdot\nabla w - \operatorname{div}\mathbb{T}(w,p)\\
&\quad = f - \delta_t - \delta\cdot\nabla\delta + \nu\operatorname{div}\mathbb{D}(\delta) = F(\delta,t) \quad \text{in } \Omega^T,\\
&\operatorname{div} w = 0 \quad \text{in } \Omega^T,\\
&w\cdot\bar n = 0 \quad \text{on } S^T,\\
&\nu\bar n\cdot\mathbb{D}(w)\cdot\bar\tau_\alpha + \gamma w\cdot\bar\tau_\alpha\\
&\quad = -\nu\bar n\cdot\mathbb{D}(\delta)\cdot\bar\tau_\alpha - \gamma\delta\cdot\bar\tau_\alpha = B_{1\alpha}(\delta),\quad \alpha=1,2, \quad \text{on } S_1^T,\\
&\bar n\cdot\mathbb{D}(w)\cdot\bar\tau_\alpha = -\bar n\cdot\mathbb{D}(\delta)\cdot\bar\tau_\alpha = B_{2\alpha}(\delta),\quad \alpha=1,2, \quad \text{on } S_2^T,\\
&w\big|_{t=0} = v(0) - \delta(0) = w(0) \quad \text{in } \Omega,
\end{aligned}
\tag{2.6}
$$

where $\operatorname{div}\delta = 0$. Moreover, we set

$$
\bar n|_{S_1} = \frac{(\varphi_{0,x_1},\varphi_{0,x_2},0)}{\sqrt{\varphi_{0,x_1}^2+\varphi_{0,x_2}^2}},\ \bar\tau_1|_{S_1} = \frac{(-\varphi_{0,x_2},\varphi_{0,x_1},0)}{\sqrt{\varphi_{0,x_1}^2+\varphi_{0,x_2}^2}},\ \bar\tau_2|_{S_1} = (0,0,1) = \bar e_3,
$$
$$
\bar n|_{S_2(-a)} = -\bar e_3,\ \bar n|_{S_2(a)} = \bar e_3,\ \bar\tau_1|_{S_2} = \bar e_1,\ \bar\tau_2|_{S_2} = \bar e_2
$$

where $\bar e_1 = (1,0,0), \bar e_2 = (0,1,0)$. For the system (2.6) we can define weak solutions and then from [RZ1] we have

Lemma 2.1. *Assume the compatibility condition (1.3). Assume that* $v(0)\in L_2(\Omega)$, $f\in L_2(0,T;L_{6/5}(\Omega))$, $d_i\in L_\infty(0,T;W_p^{s-1/p}(S_2))\bigcap L_2(0,T;W_2^{1/2}(S_2))$, $\frac{3}{p}+\frac{1}{3}\le s, p>3$ *or* $p=3, s>\frac{4}{3}$, $d_{i,t}\in L_2(0,T;W_{6/5}^{1/6}(S_2)), i=1,2$. *Then there exists a weak solution* v *to problem (1.1) such that* v *is weakly continuous with respect to* t *in* $L^2(\Omega)$ *norm and* v *converges to* v_0 *as* $t\to 0$ *strongly in* $L^2(\Omega)$ *norm. Moreover,* $v\in V_2^0(\Omega^T)$, $v\cdot\bar\tau_\alpha\in L_2(0,T;L_2(S_1)), \alpha=1,2$, *and* v *satisfies*

$$
\begin{aligned}
&\|v\|^2_{V_2^0(\Omega^t)} + \gamma\sum_{\alpha=1}^2\int_0^t \|v\cdot\bar\tau_\alpha\|^2_{L_2(S_1)} \le 2\|f\|^2_{L_2(0,t;L_{6/5}(\Omega))}\\
&+\varphi\left(\sup_{\tau\le t}\|d\|_{W_3^{s-1/p}(S_2)}\right)\left(\|d\|^2_{L_2(0,t;W_2^{1/2}(S_2))} + \|d_t\|^2_{L_2(0,t;W_{6/5}^{1/6}(S_2))}\right) + \|v(0)\|^2_{L_2(\Omega)} \equiv A^2
\end{aligned}
\tag{2.7}
$$

where φ *is a nonlinear positive increasing function of its argument and* $t\le T$.

Since we would like to have not restricted magnitudes of $v(0), f$ and d, we would need some smallness of $v_{x_3}(0)$ and f_{x_3} in L_2 norms, instead. Therefore, we set $h = v_{,x_3}$, $q = p_{,x_3}$ that are solutions to the problem: (see Lemma 3.1, [Z1].)

$$
\begin{aligned}
&h_{,t} - \operatorname{div}\mathbb{T}(h,q) = -v\cdot\nabla h - h\cdot\nabla v + g && \text{in } \Omega^T,\\
&\operatorname{div} h = 0 && \text{in } \Omega^T,\\
&\bar n\cdot h = 0,\quad \bar n\cdot\mathbb{D}(h)\cdot\bar\tau_\alpha + \gamma h\cdot\bar\tau_\alpha = 0,\quad \alpha=1,2 && \text{on } S_1^T,\\
&h_i = -d_{x_i},\quad i=1,2,\quad h_{3,x_3} = \Delta' d && \text{on } S_2^T,\\
&h\big|_{t=0} = h(0) && \text{in } \Omega.
\end{aligned}
\tag{2.8}
$$

where $g = f_{,x_3}, \Delta' = \partial^2_{x_1} + \partial^2_{x_2}$. With $\chi = (\operatorname{rot} v)_3$, we consider the problem

$$\begin{aligned} & v_{1,x_2} - v_{2,x_1} = \chi && \text{in } \Omega', \\ & v_{1,x_1} + v_{2,x_2} = -h_3 && \text{in } \Omega', \\ & v' \cdot \bar{n}' = 0 && \text{on } S'_1, \end{aligned} \tag{2.9}$$

where Ω' is a cross-section of Ω by the plane perpendicular to the x_3 axis crossing it within the interval $(-a, a)$ and $\partial\Omega' = S_1 \bigcap P \equiv S'_1$. Namely,

$$\begin{aligned} \Omega' &= \Omega \cap \{\text{plane} : x_3 = \text{const} \in (-a, a)\} \\ S'_1 &= S_1 \cap \{\text{plane} : x_3 = \text{const} \in (-a, a)\} \end{aligned}$$

and x_3, t are treated as parameters. Then, the function $\chi = (\operatorname{rot} v)_3 = v_{2,x_1} - v_{1,x_2}$ satisfies (see Lemma 3.2 in [Z1])

$$\begin{aligned} & \chi_{,t} + v \cdot \nabla\chi - h_3\chi + h_2 v_{3,x_1} - h_1 v_{3,x_2} - \nu\Delta\chi = F_3 && \text{in } \Omega^T, \\ & \chi = v_i(n_{i,x_j}\tau_{1j} + \tau_{1i,x_j}n_j) + v \cdot \bar{\tau}_1(\tau_{12,x_1} - \tau_{11,x_2}) \\ & + \frac{\gamma}{\nu} v_j \tau_{1j} \equiv \chi_* && \text{on } S_1^T, \\ & \chi_{,x_3} = 0 && \text{on } S_2^T, \\ & \chi\big|_{t=0} = \chi(0) && \text{in } \Omega, \end{aligned} \tag{2.10}$$

where $F_3 = f_{2,x_1} - f_{1,x_2}$.

3. ESTIMATES

In order to obtain the energy estimate for h – a solution to problem (2.8) we have to make the Dirichlet boundary condition on S_2^T homogeneous. For this purpose we are looking for such a function $\tilde{h}$ that

$$\begin{aligned} & \operatorname{div} \tilde{h} = 0 \quad \text{in } \Omega, \\ & \tilde{h} = 0 \quad \text{on } S_1, \\ & \tilde{h}_i = -d_{,x_i}, \ i = 1, 2 \quad \text{on } S_2, \\ & \tilde{h}_3 = 0 \quad \text{on } S_2. \end{aligned} \tag{3.1}$$

Lemma 3.1. *There exists a function* $\tilde{h}$ *satisfying (3.1) such that*

$$\begin{aligned} & \|\tilde{h}\|_{W^1_\sigma(\Omega)} \le c(\varepsilon + \rho^a)\|d_{,x'}\|_{W^1_\sigma(S_2)}, \\ & \|\tilde{h}_{,t}\|_{L_\sigma(\Omega)} \le c(\varepsilon + \rho^a)\|d_{,x't}\|_{L_\sigma(S_2)}, \end{aligned} \tag{3.2}$$

where $\sigma \in (1, \infty)$ *and* $a > 0$.

Proof. First, we construct function $\tilde{h}$. Let us define the functions

$$\begin{aligned} \bar{h}_i &= -(\eta_1 \tilde{d}_{1,x_i} + \eta_2 \tilde{d}_{2,x_i}), i = 1, 2, \\ \bar{h}_3 &= 0, \end{aligned}$$

where η_1, η_2 are smooth cut-off functions introduced before and $\tilde{d}_i$ is an extension of d_i on a neighbourhood of $S_2(a_i), i = 1, 2$. We have the compatibility condition

$$\sum_{i=1}^{2} n_i|_{S_1} \cdot d_{\alpha,x_i} = 0, \quad \alpha = 1, 2 \tag{3.3}$$

which follows from the fact that $\bar{n}|_{S_1}$ does not depend on x_3, so

$$v\cdot\bar{n}|_{S_1}=0\Rightarrow v_{,x_3}\cdot\bar{n}|_{S_1}=0\Rightarrow h\cdot\bar{n}|_{S_1}=0\Rightarrow h|_{S_2}\cdot\bar{n}|_{S_1}=0$$

and in view of $(2.8)_4$ the relation (3.3) holds.

Hence $\bar{h}$ is a solution to the problem

$$\begin{aligned}\operatorname{div}\bar{h}&=-(\eta_1\Delta'\tilde{d}_1+\eta_2\Delta'\tilde{d}_2)\quad\text{in}\ \ \Omega,\\ \bar{h}\cdot\tau_1&=-(\eta_1\tilde{d}_{1,x_i}+\eta_2\tilde{d}_{2,x_i})\tau_{1i}\quad\text{on}\ \ S_1,\\ \bar{h}\cdot\tau_2&=0\quad\text{on}\ \ S_1,\\ \bar{h}\cdot\bar{n}&=0\quad\text{on}\ \ S_1,\\ \bar{h}_i&=-d_{j,x_i},\quad i,j=1,2\quad\text{on}\ \ S_2(a_j),\\ \bar{h}_3&=0\quad\text{on}\ \ S_2.\end{aligned}$$

Next we define a function ϕ such that

$$\begin{aligned}\Delta\phi&=-(\eta_1\Delta'\tilde{d}_1+\eta_2\Delta'\tilde{d}_2),\\ \bar{n}\cdot\nabla\phi|_S&=0,\end{aligned}$$

and we are looking for a function λ and σ such that

$$\begin{aligned}-\Delta\lambda+\nabla\sigma&=0,\\ \operatorname{div}\lambda&=0,\\ \lambda\cdot\bar{\tau}_\beta&=-\bar{\tau}_\beta\cdot\nabla\phi+\bar{h}\cdot\bar{\tau}_\beta,\quad\beta=1,2,\ \ \text{on}\ S_1\\ \lambda\cdot\bar{n}&=0,\ \ \text{on}\ S_1\\ \lambda_i&=-\nabla_i\phi,\ \ i=1,2,\ \ \text{on}\ S_2\\ \lambda_3&=0\ \ \text{on}\ S_2.\end{aligned}\tag{3.4}$$

Then

$$\tilde{h}=\bar{h}-(\lambda+\nabla\phi)$$

is a solution to problem (3.1). We can get some estimates for $\tilde{h}$. We note that

$$\|\bar{h}\|_{L_p(\Omega)}\le\sum_{j=1}^2\|\eta_j\tilde{d}_{j,x'}\|_{L_p(\Omega)}\le c\rho^{1/p_0'}\sum_{j=1}^2\|\eta_j\tilde{d}_{j,x'}\|_{L_{p_0}(\Omega)},\quad\text{where}\ \ \frac{1}{p_0}+\frac{1}{p_0'}=\frac{1}{p}.$$

Let G be the Green function to the Neumann problem for ϕ. Then we have

$$\phi(x)=\int_\Omega G(x,y)\partial_{y_i}(\eta_1\tilde{d}_{1,y_i}+\eta_2\tilde{d}_{2,y_i})dy=-\int_\Omega\nabla_{y_i}G(x,y)(\eta_1\tilde{d}_{1,y_i}+\eta_2\tilde{d}_{2,y_i})dy,$$

where we used that $\bar{h}\cdot\bar{n}|_{S_1}=0$. Then we calculate

$$\nabla_x\phi(x)=-\int_\Omega\nabla_x\nabla_{y_i}G(x,y)(\eta_1\tilde{d}_{1,y_i}+\eta_2\tilde{d}_{2,y_i})dy$$

and

$$\|\nabla_x\phi\|_{L_p(\Omega)}\le\sum_{j=1}^2\|\eta_j\tilde{d}_{j,x'}\|_{L_p(\Omega)}\le c\rho^{1/p_0'}\sum_{j=1}^2\|\eta_j\tilde{d}_{j,x'}\|_{L_{p_0}(\Omega)},\qquad\text{where}\ \ \frac{1}{p_0}+\frac{1}{p_0'}=\frac{1}{p}.\tag{3.5}$$

Applying now the existence of the Green function to problem (3.4) we have

$$\lambda_i(x) = \int_{S_1} \frac{\partial G_{i\beta}}{\partial n_{S_1}} (-\bar\tau_\beta \cdot \nabla\phi + \bar h \cdot \bar\tau_\beta) dS_1 + \int_{S_2} \frac{\partial G_{ij}}{\partial n_{S_2}} (-\nabla_j \phi) dS_2.$$

In view of (3.5) we obtain

$$\|\lambda\|_{L_p(\Omega)} \le c\rho^{1/p_0'} \sum_{j=1}^{2} \|\eta_j \tilde d_{j,x'}\|_{L_{p_0}(\Omega)}. \tag{3.6}$$

Continuing the considerations and using properties of functions $\eta_i, i = 1, 2$, yields that constructed function $\tilde h$ satisfies (3.2). This concludes the proof. □

3.1. **A priori inequalities for functions k and h.** Let us introduce the new function

$$k = h - \tilde h.$$

Then k is a solution to the problem

$$\begin{aligned}
k_{,t} - \operatorname{div} \mathbb{T}(h,q) &= -v \cdot \nabla h - h \cdot \nabla v - \tilde h_{,t} + g \equiv \mathbf{g} && \text{in } \Omega^T,\\
\operatorname{div} k &= 0 && \text{in } \Omega^T,\\
\bar n \cdot k = 0, \quad \nu \bar n \cdot \mathbb{D}(h) \cdot \bar\tau_\alpha &+ \gamma h \cdot \bar\tau_\alpha = 0, \quad \alpha = 1,2 && \text{on } S_1^T,\\
k_i = 0, \quad i = 1,2, \quad h_{3,x_3} &= \Delta' d && \text{on } S_2^T,\\
k\big|_{t=0} &= h(0) - \tilde h(0) \equiv k(0) && \text{in } \Omega.
\end{aligned} \tag{3.7}$$

where $g = f_{,x_3}, \Delta' = \partial_{x_1}^2 + \partial_{x_2}^2$.We have the following Korn inequality

Lemma 3.2. [Z1] *Assume that Ω is not axially symmetric,*

$$E_\Omega(k) = \sum_{i,j=1}^{3} \int_\Omega (k_{i,x_j} + k_{j,x_i})^2 dx, \quad E_\Omega(k) + \sum_{\alpha=1}^{2} \|k \cdot \bar\tau_\alpha\|_{L_2(S_1)}^2 < \infty,$$

$$\|\Delta' d\|_{L_2(S_2)} + \|\tilde h_{3,x_3}\|_{L_2(\Omega)} + \|\tilde h_3\|_{L_2(\Omega)} < \infty.$$

Then the following inequality is valid

$$\begin{aligned}
\|k\|_{H^1(\Omega)}^2 &\le c\left(E_\Omega(k) + \sum_{\alpha=1}^{2} \gamma \|k \cdot \bar\tau_\alpha\|_{L_2(S_1)}^2\right)\\
&+ c(\|\Delta' d\|_{L_2(S_2)}^2 + \|\tilde h_{3,x_3}\|_{L_2(S_2)}^2 + \|\tilde h_3\|_{L_2(\Omega)}).
\end{aligned} \tag{3.8}$$

Let us consider the problem (3.7) in the form

$$\begin{aligned}
k_{,t} - \operatorname{div} \mathbb{T}(h,q) &= -v \cdot \nabla k - k \cdot \nabla v - v \cdot \nabla \tilde h - \tilde h \cdot \nabla v - \tilde h_{,t} + g && \text{in } \Omega^T,\\
\operatorname{div} k &= 0 && \text{in } \Omega^T,\\
\bar n \cdot k = 0, \quad \nu \bar n \cdot \mathbb{D}(h) \cdot \bar\tau_\alpha &+ \gamma h \cdot \bar\tau_\alpha = 0, \quad \alpha = 1,2 && \text{on } S_1^T,\\
k_i = 0, \quad i = 1,2, \quad h_{3,x_3} &= \Delta' d && \text{on } S_2^T,\\
k\big|_{t=0} &= h(0) - \tilde h(0) \equiv k(0) && \text{in } \Omega.
\end{aligned} \tag{3.9}$$

Using function $w = u - \delta$ we reformulate the k problem into the following system

$$
\begin{aligned}
&k_{,t} - \operatorname{div}\mathbb{T}(h,q) = -w\cdot\nabla k - k\cdot\nabla v\\
&\qquad -\delta\cdot\nabla k - v\cdot\nabla\tilde h - \tilde h\cdot\nabla v - \tilde h_{,t} + g \equiv G && \text{in } \Omega^T,\\
&\operatorname{div} k = 0 && \text{in } \Omega^T,\\
&\bar n\cdot k = 0,\quad \nu\bar n\cdot\mathbb{D}(h)\cdot\bar\tau_\alpha + \gamma h\cdot\bar\tau_\alpha = 0,\quad \alpha = 1,2 && \text{on } S_1^T,\\
&k_i = 0,\quad i = 1,2,\quad h_{3,x_3} = \Delta' d && \text{on } S_2^T,\\
&k\big|_{t=0} = k(0) && \text{in } \Omega.
\end{aligned}
\tag{3.10}
$$

Projecting $\operatorname{div} k$ on S_2 we see that

$$\operatorname{div} k|_{S_2} = k_{3,x_3}|_{S_2} = 0$$

Then the second condition in $(3.10)_4$ takes the form

$$\tilde h_{3,x_3} = \Delta' d \quad \text{on} \quad S_2$$

Lemma 3.3. *Let*

$$
\begin{aligned}
\Lambda &= D_1 + D_2 + \|f_3\|^2_{L_2(0,t;L_{4/3}(S_2))}\\
&\quad + \|g\|^2_{L_2(0,t;L_{6/5}(\Omega))} + \|h(0)\|^2_{L_2(\Omega)},\\
D_1 &= \|d_t\|^2_{L_2(0,t;H^1(S_2))} + \|d_{,x'}\|^2_{L_2(0,t;H^1(S_2))}\\
D_2 &= \|d_{,x'}\|^2_{L_\infty(0,t;H^1(S_2))},
\end{aligned}
$$

and

$$H_1 = \|h\|^2_{L_2(\Omega^t)} + \|h\|^2_{L_\infty(0,t;L_3(\Omega))}. \tag{3.11}$$

Then the solution k to the equivalent problems (3.7, 3.9,3.10) satisfies

$$
\begin{aligned}
&\|k\|^2_{V_2^0(\Omega^t)} + \sum_{\alpha=1}^2 \|k\cdot\bar\tau_\alpha\|^2_{L_2(S_1^t)}\\
&\le c(\|d\|^6_{L_\infty(0,t;L_3(S_2))} + A^2 + 1)(\Lambda^2 + H_1^2)
\end{aligned}
\tag{3.12}
$$

Proof. We shall obtain the energy type estimate for solutions to problem (3.10). Multiplying $(3.10)_1$ by k and integrating over Ω yields

$$
\begin{aligned}
&\frac{1}{2}\frac{d}{dt}\int_\Omega k^2dx - \int_\Omega \operatorname{div}\mathbb{T}(h,q)kdx = -\int_\Omega w\cdot\nabla kkdx - \int_\Omega k\nabla v\cdot kdx\\
&-\int_\Omega \delta\cdot\nabla kkdx - \int_\Omega v\cdot\nabla\tilde hkdx - \int_\Omega \tilde h\cdot\nabla vkdx - \int_\Omega \tilde h_tkdx + \int_\Omega gkdx.
\end{aligned}
\tag{3.13}
$$

Now we examine the particular terms in (3.13). Integrating by parts the second term on the l.h.s takes the form

$$-\int_{S_1}\bar n\cdot\mathbb{T}(h,q)\cdot kdS_1 - \int_{S_2}\bar n\cdot\mathbb{T}(h,q)\cdot kdS_2 + \frac{\gamma}{2}\int_\Omega \mathbb{D}(h)\mathbb{D}(k)dx \equiv I_1 + I_2 + I_3,$$

where

$$
\begin{aligned}
I_1 &= -\int_{S_1} \bar{n}\cdot\mathbb{T}(h,q)\cdot\bar{\tau}_\alpha k\cdot\bar{\tau}_\alpha dS_1 = \gamma\int_{S_1} h\cdot\bar{\tau}_\alpha k\cdot\bar{\tau}_\alpha dS_1\\
&= \gamma\|k\cdot\bar{\tau}_\alpha\|^2_{L_2(S_1)} + \gamma\int_{S_1}\tilde{h}\cdot\bar{\tau}_\alpha k\cdot\bar{\tau}_\alpha dS_1,
\end{aligned}
$$

$$
\begin{aligned}
I_2 = -\int_{S_2}\mathbb{T}_{nn}(h,q)k_n dS_2 &= -\int_{S_2}(2\nu h_{3,x_3}-q)k_n dS_2 = -\int_{S_2}(2\nu\tilde{h}_{3,x_3}-q)k_n dS_2\\
&= -2\nu\int_{S_2}\tilde{h}_{3,x_3}k_n dS_2 - \int_{S_2(-a)} qk_3 dS_2 + \int_{S_2(a)} qk_3 dS_2
\end{aligned}
$$

To examine the last integral we use the third component of $(1.1)_1$ projected on S_2

$$
d_t + v\cdot\nabla v_3 - \nu\Delta' d - \nu\tilde{h}_{3,x_3} - f_3 = -q \tag{3.14}
$$

Using this relation in the last term of I_2 we obtain

$$
\begin{aligned}
\int_{S_2} qk_3 dS_2 = \int_{S_2}(-d_t + \nu\Delta' d + \nu\tilde{h}_{3,x_3} + f_3)k_3 dS_2 - \int_{S_2} v'\nabla' dk_3 dS_2\\
+\int_{S_2(-a)} dk_3^2 dS_2 - \int_{S_2(a)} dk_3^2 dS_2
\end{aligned}
$$

where

$$
d|_{S_2(-a)} = -d_1, d_1 > 0, d|_{S_2(a)} = d_2 > 0
$$

and in the first two terms we do not distinguish dependence on $S_2(-a)$ and $S_2(a)$ because it does not have any influence on estimations.

The first term in the above expression we estimate by

$$
\varepsilon_1\|k_3\|^2_{L_4(S_2)} + c(1/\varepsilon_1)\left(\|d_t\|^2_{L_{4/3}(S_2)} + \|\Delta' d\|^2_{L_{4/3}(S_2)} + \|f_3\|^2_{L_{4/3}(S_2)}\right),
$$

the second term by

$$
\varepsilon_2\|k_3\|^2_{L_4(S_2)} + c(1/\varepsilon_2)\|v'\|^2_{L_4(S_2)}\|\nabla' d\|^2_{L_2(S_2)}
$$

and the last one as follows

$$
\begin{aligned}
\int_{S_2(-a)} d_1 k_3^2 dS_2 \le \|d\|_{L_3(S_2)}\|k_3\|^2_{L_3(S_2)} \le \left(\varepsilon^{1/6}\|\nabla k_3\|^2_{L_2(\Omega)} + c\varepsilon^{-5/6}\|k_3\|^2_{L_2(\Omega)}\right)\cdot\|d\|_{L_3(S_2)}\\
\le \varepsilon_2\|\nabla k_3\|^2_{L_2(\Omega)} + c\varepsilon_2^{-5/6}\|d\|^6_{L_3(S_2)}\cdot(\|h_3\|^2_{L_2(\Omega)} + \|\tilde{h}_3\|^2_{L_2(\Omega)}).
\end{aligned}
$$

Using the above estimates of the second term on the l.h.s. of (3.13) and denoting the r.h.s. of (3.10) by G we obtain

$$
\begin{aligned}
\frac{1}{2}\frac{d}{dt}\int_\Omega k^2 dx + \frac{\nu}{2}\int_\Omega|\mathbb{D}(k)|^2 dx + \frac{\gamma}{2}\int_{S_1}|k\cdot\bar{\tau}_\alpha|^2 dS_1 \le \frac{\nu}{4}\|k\|^2_{H^1(\Omega)} + \frac{\gamma}{2}\int_{S_1}|\tilde{h}\cdot\bar{\tau}_\alpha|^2 dS_1\\
+c\int_\Omega|\mathbb{D}(\tilde{h})|^2 dx + c(\|d_t\|^2_{L_{4/3}(S_2)} + \|\Delta' d\|^2_{L_{4/3}(S_2)} + \|f_3\|^2_{L_{4/3}(S_2)})\\
+\|d\|^6_{L_3(S_2)}\cdot(\|h_3\|^2_{L_2(\Omega)} + \|\tilde{h}_3\|^2_{L_2(\Omega)}) + |\int_\Omega Ghdx|
\end{aligned}
$$

We apply the Korn inequality (3.8) to conclude

$$\begin{aligned}&\frac{d}{dt}\int_\Omega k^2dx+\nu\|k\|^2_{H^1(\Omega)}+\gamma\|k\cdot\bar\tau_\alpha\|^2_{L_2(S_1)}\\ (3.15)\quad&\le c(\|d_t\|^2_{L_{4/3}(S_2)}+\|\Delta'd\|^2_{L_{4/3}(S_2)}+\|f_3\|^2_{L_{4/3}(S_2)}+\|\tilde h\|^2_{H^1(\Omega)})\\ &+\|d\|^6_{L_3(S_2)}\cdot(\|h_3\|^2_{L_2(\Omega)}+\|\tilde h_3\|^2_{L_2(\Omega)})+|\int_\Omega Ghdx|\end{aligned}$$

Now we shall examine the last term on the r.h.s. of the inequality (3.15). To this end we use the r.h.s. of (3.13). The first term on the r.h.s. of (3.13) vanishes. The second term can be estimated either by

$$\int_\Omega k^2|\nabla v|dx\le\varepsilon_3\|k\|^2_{L_6(\Omega)}+c(1/\varepsilon_3)\|\nabla v\|^2_{L_2(\Omega)}\|k\|^2_{L_3(\Omega)}$$

or by

$$\int_\Omega k^2|\nabla v|dx\le\varepsilon_3\|k\|^2_{L_6(\Omega)}+c(1/\varepsilon_3)\|\nabla v\|^2_{L_3(\Omega)}\|k\|^2_{L_2(\Omega)}.$$

To examine the third term on the r.h.s. of (3.13) we express it in the form

$$\int_\Omega b\nabla kkdx+\int_\Omega\nabla\varphi\nabla kkdx\equiv I_1+I_2$$

In view of [RZ1] we have

$$|I_1|\le c\rho^{1/6}\|\tilde d\|_{H^1(\Omega)}\|k\|^2_{H^1(\Omega)}$$

and

$$|I_2|\le c(\varepsilon_4\rho^{\mu-2/3}\sup_{x_3}\|\tilde d_3\|_{L_3(S_2)}+\rho^{\mu+1/3}\sup_{x_3}\|\tilde d_{,x_3}\|_{L_3(S_2)})\|k\|^2_{H^1(\Omega)}$$

The fourth term we estimate by

$$\varepsilon_5\|k\|^2_{L_6(\Omega)}+c(1/\varepsilon_5)\|v\|^2_{L_6(\Omega)}\|\nabla\tilde h\|^2_{L_{3/2}(\Omega)}$$

The fifth term we treat as follows

$$\varepsilon_6\|k\|^2_{L_6(\Omega)}+c(1/\varepsilon_6)\|\tilde h\|^2_{L_3(\Omega)}\|\nabla v\|^2_{L_2(\Omega)},$$

the sixth by

$$\varepsilon_7\|k\|^2_{L_6(\Omega)}+c(1/\varepsilon_7)\|\tilde h_t\|^2_{L_{6/5}(\Omega)}$$

and finally the last one by

$$\varepsilon_8\|k\|^2_{L_6(\Omega)}+c(1/\varepsilon_8)\|g\|^2_{L_{6/5}(\Omega)}$$

Using the above considerations in (3.13) and assuming that $\varepsilon_1-\varepsilon_8$ are sufficiently small we obtain

$$\begin{aligned}&\frac{d}{dt}\int_\Omega k^2dx+\nu\|k\|^2_{H^1(\Omega)}+\gamma\sum_{\alpha=1}^2\|k\cdot\bar\tau_\alpha\|^2_{L_2(S_1)}\\ (3.16)\quad&\le c\Big[\|\Delta'd\|^2_{L_2(S_2)}+\|d_t\|^2_{L_{4/3}(S_2)}+\|f_3\|^2_{L_{4/3}(S_2)}+\|\tilde h\|^2_{H^1(\Omega)}\\ &+\|d\|^6_{L_3(S_2)}(\|h_3\|^2_{L_2(\Omega)}+\|\tilde h_3\|^2_{L_2(\Omega)})+\|\nabla v\|^2_{L_2(\Omega)}(\|h_3\|^2_{L_3(\Omega)}+\|\tilde h_3\|^2_{L_3(\Omega)})\\ &+\|v\|^2_{L_6(\Omega)}\|\nabla\tilde h\|^2_{L_{3/2}(\Omega)}+\|\tilde h\|^2_{L_3(\Omega)}\|\nabla v\|^2_{L_2(\Omega)}+\|h_t\|^2_{L_{6/5}(\Omega)}+\|g\|^2_{L_{6/5}(\Omega)}\Big].\end{aligned}$$

Integrating (3.16) with respect to time and using the energy type bound of the form

$$\|v\|_{V_2^0(\Omega^t)} \le A, \quad t \le T,$$

where A is defined in Lemma 2.1 we get

$$\begin{aligned}
\|k\|^2_{V_2^0(\Omega^t)} + \gamma\sum_{\alpha=1}^{2}\|k\cdot\bar\tau_\alpha\|^2_{L_2(S_1^t)} &\le c\Big[\|\Delta' d\|^2_{L_2(0,t;L_2(S_2))} + \|d_t\|^2_{L_2(0,t;L_{4/3}(S_2))} + \|f_3\|^2_{L_2(0,t;L_{4/3}(S_2))}\\
&\quad + \|\tilde h\|^2_{H^1(\Omega^t)} + \|d\|^6_{L_\infty(0,t;L_3(S_2))}\big(\|h_3\|^2_{L_2(\Omega^t)} + \|\tilde h_3\|^2_{L_2(\Omega^t)}\big)\Big]\\
&\quad + \sup_\tau A^2\|h\|^2_{L_3(\Omega)} + \sup_\tau A^2\|\nabla\tilde h\|^2_{L_{3/2}(\Omega)} + \sup_\tau A^2\|\tilde h\|^2_{L_3(\Omega)} + \|\tilde h_t\|^2_{L_2(\Omega^t)}\\
&\quad + \|g\|^2_{L_2(0,t;L_{6/5}(\Omega))} + \|h(0)\|^2_{L_2(\Omega)} + \|\tilde h(0)\|^2_{L_2(\Omega)}.
\end{aligned}\tag{3.17}$$

In view of Lemma 3.1 we have

$$\begin{aligned}
\|k\|^2_{V_2^0(\Omega^t)} + \gamma\sum_{\alpha=1}^{2}\|k\cdot\bar\tau_\alpha\|^2_{L_2(S_1^t)} &\le c\Big[\|d_{,x'}\|^2_{L_2(0,t;H^1(S_2))} + \|d_t\|^2_{L_2(0,t;L_{4/3}(S_2))}\\
&\quad + \|f_3\|^2_{L_2(0,t;L_{4/3}(S_2))} + \|d\|^6_{L_\infty(0,t;L_3(S_2))}\|h_3\|^2_{L_2(\Omega^t)}\Big]\\
&\quad + A^2\big(\sup_\tau\|h\|^2_{L_3(\Omega)} + A^2\|d_{,x'}\|^2_{L_\infty(0,t;H^1(S_2))}\big) + A^2\sup_\tau\|d_{,x't}\|^2_{L_2(S_2)}\\
&\quad + \|g\|^2_{L_2(0,t;L_{6/5}(\Omega))} + \|h(0)\|^2_{L_2(\Omega)}.
\end{aligned}\tag{3.18}$$

We set

$$D_1 = \|d_t\|^2_{L_2(0,t;H^1(S_2))} + \|d_{,x'}\|^2_{L_2(0,t;H^1(S_2))}$$

$$D_2 = \|d_{,x'}\|^2_{L_\infty(0,t;H^1(S_2))}$$

Then (3.18) takes the form

$$\begin{aligned}
\|k\|^2_{V_2^0(\Omega^t)} + \gamma\sum_{\alpha=1}^{2}\|k\cdot\bar\tau_\alpha\|^2_{L_2(S_1^t)} &\le c\Big[D_1^2 + \|f_3\|^2_{L_2(0,t;L_{4/3}(S_2))}\\
&\quad + (\|d\|^6_{L_\infty(0,t;L_3(S_2))} + A^2 + 1)\Big(D_2^2 + \|h\|^2_{L_2(\Omega^t)} + \|h\|^2_{L_\infty(0,t;L_3(\Omega))}\Big)\\
&\quad + \|g\|^2_{L_2(0,t;L_{6/5}(\Omega))} + \|h(0)\|^2_{L_2(\Omega)}\Big].
\end{aligned}\tag{3.19}$$

Therefore, defining quantities Λ, D_1, D_2, H_1 as in (1.7) and (3.11) we can formulate the following compact version of the last inequality.

$$\begin{aligned}
&\|k\|^2_{V_2^0(\Omega^t)} + \sum_{\alpha=1}^{2}\|k\cdot\bar\tau_\alpha\|^2_{L_2(S_1^t)}\\
&\le c(\|d\|^6_{L_\infty(0,t;L_3(S_2))} + A^2 + 1)(\Lambda^2 + H_1^2)
\end{aligned}$$

and this concludes the proof. □

Consequently, we can show the following inequality for h

Lemma 3.4. *Let D_0, Λ are finite. Then a solutions h of (2.8) satisfies*

$$\|h\|^2_{V_2^0(\Omega^t)} + \sum_{\alpha=1}^{2}\|h\cdot\bar\tau_\alpha\|^2_{L_2(S_1^t)} \le cD_0(\Lambda^2 + H_1^2)\tag{3.20}$$

where

$$D_0 = \|d\|^6_{L_\infty(0,t;L_3(S_2))} + A^2 + 1$$

Proof. Using (3.12), $h = k + \tilde{h}$, and the estimate

$$\|\tilde{h}\|^2_{V_2^0(\Omega^t)} + \sum_{\alpha=1}^2 \|\tilde{h} \cdot \bar{\tau}_\alpha\|^2_{L_2(S_1^t)} \le c(D_1^2 + D_2^2)$$

we conclude the result. □

Remark 3.5. *Since we want to admit arbitrary large flux, the inequality above does not involve any small parameter that we could use in the existence proof. Thus, we will need some more refined relation and this can be achieved by improving the regularity through the system for vorticity component χ.*

3.2. **A priori estimates for vorticity component χ.** We need to examine solutions of problem (2.9) and consequently, (2.10). We have to deal with possibly non-zero boundary condition of function χ on S_1^T which we set as χ_*. To this end, let us introduce a function $\tilde{\chi}$ as a solution to the following problem

$$\begin{aligned} \tilde{\chi}_{,t} - \nu\Delta\tilde{\chi} &= 0 \quad \text{in } \Omega^T, \\ \tilde{\chi} &= \chi_* \quad \text{on } S_1^T, \\ \tilde{\chi}_{,x_3} &= 0, \quad \text{on } S_2^T, \\ \tilde{\chi}|_{t=0} &= \chi_0 \quad \text{in } \Omega. \end{aligned} \tag{3.21}$$

To show the existence of such function we need the following compatibility conditions

$$\begin{aligned} &\chi_{*,x_3} = 0, \text{ on } \bar{S}_1 \cap \bar{S}_2 \\ &\chi_0|_{S_1} = \chi_*|_{t=0}, \\ &\chi_{0,x_3} = 0 \text{ on } S_2. \end{aligned} \tag{3.22}$$

where

$$\chi_{*,x_3} = -\sum_{i,j=1}^2 \Big[d_{,x_i}(n_{i,x_j}\tau_{1j} + \tau_{1i,x_j}n_j) + \frac{\gamma}{\nu} d_{,x_j}\tau_{1j} + d_{x_i}\tau_{1i}(\tau_{12,x_1} - \tau_{11,x_2}) \Big]$$

We note that χ_0 depends of v in a similar way as χ_*.

Then, we consider the new function $\chi' = \chi - \tilde{\chi}$ which is a solution to the following problem

$$\begin{aligned} &\chi'_{,t} + v \cdot \nabla\chi' - h_3\chi' + h_2 v_{3,x_1} - h_1 v_{3,x_2} - \nu\Delta\chi' \\ &= F_3 - v \cdot \nabla\tilde{\chi} + h_3\tilde{\chi} && \text{in } \Omega^T, \\ &\chi' = 0 && \text{on } S_1^T, \\ &\chi'_{,x_3} = 0 && \text{on } S_2^T, \\ &\chi'\big|_{t=0} = \chi(0) && \text{in } \Omega. \end{aligned} \tag{3.23}$$

Lemma 3.6. *Assume that $h \in L_\infty(0,t;L_3(\Omega))$, $v' \in L_\infty(0,t;H^1(\Omega)) \cap L_2(0,t;H^2(\Omega)) \cap L_2(\Omega;H^{1/2}(0,t))$, $\chi(0) \in L_2(\Omega)$, $F_3 \in L_2(0,t;L_{6/5}(\Omega))$. Then solutions to problem*

(2.10) satisfy the inequality

$$
\begin{aligned}
\|\chi\|_{V_2^0(\Omega^t)} &\le \varphi(c_2,A)\sup_t\|h(t)\|_{L_3(\Omega)} + \frac{1}{\varepsilon}\varphi(c_2,c_3,A)\sup_t\|h(t)\|^2_{L_3(\Omega)}\\
&+\frac{1}{\varepsilon}\varphi(c_2,c_3,A) + \frac{\gamma c_2}{\nu}\|F_3\|_{L_2(0,t;H^2(\Omega))} + \varepsilon(\|v'\|_{L_\infty(0,t;H^1(\Omega))}\\
&+\|v'\|_{L_2(0,t;H^2(\Omega))}) + c_3\|v'\|_{L_2(\Omega;H^{1/2}(0,t))} + \|\chi(0)\|_{L_2(\Omega)}
\end{aligned}
\tag{3.24}
$$

$\equiv c(H_1+H_1^2) + c\|v'\|_{L_2(\Omega;H^{1/2}(0,t))} + \varepsilon(\|v'\|_{L_\infty(0,t;H^1(\Omega))} + \|v'\|_{L_2(0,t;H^2(\Omega))}) + c\Lambda_2,$

where $\varepsilon\in(0,1), a>0, t\in(0,T), c_3$ *is some constant,*

$$\Lambda_2 = \|F_3\|_{L_2(0,t;H^2(\Omega))} + \|\chi(0)\|_{L_2(\Omega)} + 1$$

and φ *is a generic function which changes its form from formula to formula.*

Proof. We use the first equation of the system (3.23): we multiply it by χ', integrate over Ω and use boundary conditions that yields

$$
\begin{aligned}
\frac{1}{2}\frac{d}{dt}\|\chi'\|^2_{L_2(\Omega)} + \nu\|\nabla\chi'\|^2_{L_2(\Omega)} &= \int_\Omega h_3\chi'^2dx - \int_\Omega(h_2v_{3,x_1} - h_1v_{3,x_2})\chi'dx + \int_\Omega F_3\chi'dx\\
&- \int_\Omega v\cdot\nabla\tilde\chi\chi'dx + \int_\Omega h_3\tilde\chi\chi'dx
\end{aligned}
\tag{3.25}
$$

The first term on the r.h.s. of (3.25) can be bounded by

$$
\begin{aligned}
&|\int_\Omega h_3\chi'^2dx| = |\int_\Omega h_3\chi'(\chi-\tilde\chi)dx| \le |\int_\Omega h_3\chi'(\chi-\tilde\chi)dx| \le |\int_\Omega h_3\chi'\chi dx| + |\int_\Omega h_3\chi'\tilde\chi dx|\\
&\quad\le \frac{\varepsilon_1}{4}\|\chi'\|^2_{L_6(\Omega)} + \frac{1}{\varepsilon_1}\|h_3\|^2_{L_3(\Omega)}\|\chi\|^2_{L_2(\Omega)} + \frac{\varepsilon_1}{4}\|\chi'\|^2_{L_6(\Omega)} + \frac{1}{\varepsilon_1}\|h_3\|^2_{L_2(\Omega)}\|\tilde\chi\|^2_{L_3(\Omega)}
\end{aligned}
$$

the second as follows

$$\frac{\varepsilon_2}{2}\|\chi'\|^2_{L_6(\Omega)} + \frac{1}{2\varepsilon_2}\|h\|^2_{L_3(\Omega)}\|v_{3,x'}\|^2_{L_2(\Omega)},$$

and the third one:

$$\frac{\varepsilon_3}{2}\|\chi'\|^2_{L_6(\Omega)} + \frac{1}{2\varepsilon_3}\|F_3\|^2_{L_{6/5}(\Omega)}.$$

The fourth term on the r.h.s. of (3.25) we express in the form

$$\int_\Omega v\cdot\nabla\chi'\tilde\chi dx$$

and estimate as follows

$$\frac{\varepsilon_4}{2}\|\nabla\chi'\|^2_{L_2(\Omega)} + \frac{1}{2\varepsilon_4}\|v\|^2_{L_6(\Omega)}\|\chi'\|^2_{L_3(\Omega)}.$$

Finally, the last term on the r.h.s. of (3.25) is bounded by

$$\frac{\varepsilon_5}{2}\|\chi'\|^2_{L_6(\Omega)} + \frac{1}{2\varepsilon_5}\|\chi'\|^2_{L_3(\Omega)}.$$

Using the above estimates in (3.25), setting $\varepsilon_1=\varepsilon_2=\varepsilon_3=\varepsilon$, $\varepsilon_4=\frac{\nu}{2}$, $\varepsilon=\frac{\nu}{8c_2}$ we obtain

$$
\begin{aligned}
\frac{d}{dt}\|\chi'\|^2_{L_2(\Omega)} + \nu\|\nabla\chi'\|^2_{L_2(\Omega)} &\le \frac{8c_2}{\nu}(\|\chi\|^2_{L_2(\Omega)} + \|v_{3,x'}\|^2_{L_2(\Omega)})\|h\|^2_{L_3(\Omega)}\\
&+\left(\frac{2}{\nu}\|v\|^2_{L_6(\Omega)} + \frac{16c_2}{\nu}\|h\|^2_{L_2(\Omega)}\right)\|\tilde\chi\|^2_{L_3(\Omega)} + \frac{8c_2}{\nu}\|F_3\|^2_{L_{6/5}(\Omega)}.
\end{aligned}
$$

Integrating the above inequality with respect to time and using estimate for the weak solutions we can find the following

$$\|\chi'\|^2_{L_2(\Omega)} + \nu\|\nabla\chi'\|^2_{L_2(\Omega^T)} \le \frac{16c_2A^2}{\nu}\sup_t\|h(t)\|^2_{L_3(\Omega)} + \frac{16c_2+2}{\nu}A^2\sup_t\|\tilde\chi\|^2_{L_3(\Omega)} + \frac{8c_2}{\nu}\|F_3\|^2_{L_2(0,T;L_{6/5}(\Omega))} + \|\chi(0)\|^2_{L_2(\Omega)}.$$

Since $\chi = \chi' + \tilde\chi$ we obtain

$$\|\chi\|^2_{V_2^0(\Omega^T)} \le \frac{16c_2A^2}{\nu}\sup_t\|h(t)\|^2_{L_3(\Omega)} + \frac{16c_2+2}{\nu}A^2\sup_t\|\tilde\chi\|^2_{L_3(\Omega)} + \frac{8c_2}{\nu}\|F_3\|^2_{L_2(0,T;L_{6/5}(\Omega))} + \|\chi(0)\|^2_{L_2(\Omega)}. \tag{3.26}$$

In view of Lemma and some interpolation inequalities we have

$$\begin{aligned}
\|\tilde\chi\|_{L_2(0,T;L_2(\Omega))} &\le c_1'\|v'\|_{L_2(0,T;L_2(S_1))} \le \varepsilon\|v'\|_{L_2(0,T;H^1(\Omega))} + c_2'\varepsilon^{-1}A,\\
\|\tilde\chi\|_{L_\infty(0,T;L_3(\Omega))} &\le c_3'\|v'\|_{L_\infty(0,T;L_3(S_1))} \le \varepsilon^{1/6}\|v'\|_{L_\infty(0,T;H^1(\Omega))} + c_4'\varepsilon^{-5/6}A,\\
\|\tilde\chi\|_{L_\infty(0,T;L_2(\Omega))} &\le c_5'\|v'\|_{L_\infty(0,T;L_2(S_1))}\\
&\le \varepsilon^{1/2}\|v'\|_{L_\infty(0,T;H^1(\Omega))} + c_6'\varepsilon^{-1/2}A,\\
\|\nabla\tilde\chi\|_{L_2(0,T;L_2(\Omega))} &\le c_7'\|v'\|_{W_2^{1/2,1/4}(S_1^T)} \le c_8'\|v'\|_{W_2^{1,1/2}(\Omega^T)}\\
&= c_8'(\|v'\|_{L_2(0,T;H^1(\Omega))} + \|v'\|_{L_2(\Omega;H^{1/2}(0,T))})\\
&\le \varepsilon^{1/2}\|v'\|_{L_2(0,T;H^2(\Omega))} + \varepsilon^{-1/2}c_9'A + c_8'\|v'\|_{L_2(\Omega;H^{1/2}(0,T))}
\end{aligned} \tag{3.27}$$

where $c_1',\dots,c_9'$ are constants from corresponding theorems and of imbedding. Let $c_i' \le c_3'$, $i = 1,\dots,9$. Then we use (3.26) and (3.27) to obtain (3.24). This concludes the proof. □

3.3. **(rot, div) problem and inequalities relating v and h.** Let us consider the elliptic (rot, div) problem (2.9), i.e.

$$\begin{aligned}
v_{1,x_2} - v_{2,x_1} &= \chi && \text{in } \Omega',\\
v_{1,x_1} + v_{2,x_2} &= -h_3 && \text{in } \Omega',\\
v'\cdot\bar n' &= 0 && \text{on } S_1',
\end{aligned}$$

For solutions v' of this system, in view of inequalities for h and χ, i.e. (3.20) and (3.24), we derive at

$$\begin{aligned}
\|v'\|^2_{V_2^1(\Omega^T)} &\le cD_0(\Lambda + H_1)\\
&\quad + c\|v'\|_{L_2(\Omega;H^{1/2}(0,t))} + c(H_1 + H_1^2) + c\Lambda_2
\end{aligned} \tag{3.28}$$

Let us consider problem (1.1) written in the form of Stokes-type system

$$\begin{aligned}
&v_{,t} - \operatorname{div}\mathbb{T}(v,p) = -v'\cdot\nabla v - wh + f, && \text{in } \Omega^T\\
&\operatorname{div} v = 0, && \text{in } \Omega^T\\
&v\cdot\bar n = 0,\ \bar n\cdot\mathbb{D}(v)\cdot\bar\tau_\alpha + \gamma v\cdot\bar\tau_\alpha = 0,\quad \alpha = 1,2, && \text{on } S_1^T\\
&v\cdot\bar n = d,\ \bar n\cdot\mathbb{D}(v)\cdot\bar\tau_\alpha = 0,\quad \alpha = 1,2, && \text{on } S_1^T\\
&v|_{t=0} = v(0), && \text{in } \Omega.
\end{aligned} \tag{3.29}$$

This yields the next result

Lemma 3.7. *Let the assumptions of Lemmas 3.1-3.5 be satisfied. Let* $h \in L_{\frac{10}{3}}(\Omega^T)$, $f \in L_2(\Omega^T), v(0) \in H^1(\Omega)$. *Then for solutions of (3.29) we obtain the inequality*

$$\begin{aligned}\|v\|_{W_2^{2,1}(\Omega^t)} + \|\nabla p\|_{L_2(\Omega^t)} &\le cD_0^2(H_1^2 + \Lambda^2) + c(H_1 + H_1^2)\\ &\quad + c\Lambda_2^2 + c\Lambda_3^2 + c\Lambda_4, \quad t \le T,\end{aligned} \tag{3.30}$$

where Λ_3, Λ_4 *are defined by (1.5).*

Proof. In view of [Z2], Lemma 3.7 we have that

$$\|v'\|_{L_{10}(\Omega^T)} \le c\|v'\|_{V_2^1(\Omega^T)}.$$

Hence

$$\begin{aligned}\|v'\nabla v\|_{L_{5/3}(\Omega^T)} &\le cA\|v'\|_{V_2^1(\Omega^T)},\\ \|wh\|_{L_{5/3}(\Omega^T)} &\le cA\|h\|_{L_{10/3}(\Omega^T)}.\end{aligned}$$

In view of the above estimates we have

$$\begin{aligned}\|v\|_{W^{2,1}_{5/3}(\Omega^T)} &\le cA(\|v'\|_{V_2^1(\Omega^T)} + \|h\|_{L_{10/3}(\Omega^T)})\\ &\quad + c(\|f\|_{L_{5/3}(\Omega^T)} + \|v(0)\|_{W^{4/5}_{5/3}(\Omega)}).\end{aligned} \tag{3.31}$$

We calculate

$$\begin{aligned}\|v'\nabla v\|_{L_2(\Omega^T)} &\le \|v'\|_{L_{10}(\Omega^T)}\|\nabla v\|_{L_{5/2}(\Omega^T)}\\ &\le \|v'\|_{V_2^1(\Omega^T)}\|v\|_{W^{2,1}_{5/3}(\Omega^T)},\\ \|wh\|_{L_2(\Omega^T)} &\le \|w\|_{L_5(\Omega^T)}\|h\|_{L_{10/3}(\Omega^T)}\\ &\le c\|v\|_{W^{2,1}_{5/3}(\Omega^T)}\|h\|_{L_{10/3}(\Omega^T)}.\end{aligned}$$

Applying (3.28) in (3.31) and using the interpolation

$$\|v'\|_{L_2(0,T;H^{1/2}(\Omega))} \le \varepsilon\|v'\|_{W^{2,1}_{5/3}(\Omega^T)} + c(1/\varepsilon)A,$$

we obtain

$$\|v\|_{W^{2,1}_{5/3}(\Omega^T)} \le cD_0(H_1 + \Lambda) + c(H_1 + H_1^2) + c\Lambda_2 + c\Lambda_3, \tag{3.32}$$

and

$$\|v\|_{W_2^{2,1}(\Omega^T)} \le cD_0^2(H_1^2 + \Lambda^2) + c(H_1 + H_1^2) + c\Lambda_2^2 + c\Lambda_3^2 + c\Lambda_4 \tag{3.33}$$

where

$$\begin{aligned}\Lambda_3 &= \|f\|_{L_{5/3}(\Omega^T)} + \|v(0)\|_{W^{4/5}_{5/3}(\Omega)},\\ \Lambda_4 &= \|f\|_{L_2(\Omega^T)} + \|v(0)\|_{H^1(\Omega)}.\end{aligned}$$

Applying the above estimates we conclude for solutions to problem (3.29) the inequality (3.30). This yields the thesis of the lemma. ☐

Next, we need to find bound for h in terms of v with some small parameter. We prove

Lemma 3.8. *Let the assumptions of Lemmas 3.1-3.5 be satisfied. Let* $\nabla v \in L_3(\Omega)$ *and* D_0, Λ *are finite. Then a solution* h *of (2.8) satisfies*

$$\|h\|^2_{V_2^0(\Omega^t)} \le D_0 \exp\left[\int_0^t (\|d\|^6_{L_3(S_2)} + \|\nabla v\|^2_{L_3(\Omega)})dt\right]\Lambda^2. \tag{3.34}$$

Proof. To this end we consider again the inequality

$$\begin{aligned}\frac{d}{dt}\int_\Omega k^2dx+\nu\|k\|^2_{H^1(\Omega)}+\gamma\|k\cdot\bar\tau_\alpha\|^2_{L_2(S_1)}&\\ \le c(\|d_t\|^2_{L_{4/3}(S_2)}+\|\Delta'd\|^2_{L_{4/3}(S_2)}+\|f_3\|^2_{L_{4/3}(S_2)}+\|\tilde h\|^2_{H^1(\Omega)})&\\ +\|d\|^6_{L_3(S_2)}\|k\|^2_{L_2(\Omega)}+|\int_\Omega Ghdx|&\end{aligned}$$

and to examine the last term on the r.h.s. we use the (3.13) and the relation

$$\int_\Omega|\nabla v|k^2dx\le\varepsilon_3\|k\|^2_{L_6(\Omega)}+c(1/\varepsilon_3)\|\nabla v\|^2_{L_3(\Omega)}\|k\|^2_{L_2(\Omega)}.$$

We deal with inequalities above and through such modifications, we have

$$\begin{aligned}\|k\|^2_{V_2^0(\Omega^t)}+\gamma\sum_{\alpha=1}^2\|k\cdot\bar\tau_\alpha\|^2_{L_2(S_1^t)}\le c\Big[\|\nabla v\|^2_{L_2(0,t;L_3(\Omega))}+\|d\|^6_{L_\infty(0,t;L_3(S_2))}\Big]\|k\|^2_{L_2(\Omega^t)}&\\ +\|d_{,x'}\|^2_{L_2(0,t;H^1(S_2))}+\|d_t\|^2_{L_2(0,t;L_{4/3}(S_2))}+\|f_3\|^2_{L_2(0,t;L_{4/3}(S_2))}&\\ +A^2\|d_{,x'}\|^2_{L_\infty(0,t;H^1(S_2))})+A^2\sup_\tau\|d_{,x't}\|^2_{L_2(S_2)}+\|g\|^2_{L_2(0,t;L_{6/5}(\Omega))}+\|h(0)\|^2_{L_2(\Omega)}.&\end{aligned}$$

Then, instead of (3.20) we can obtain

$$\|k\|^2_{V_2^0(\Omega^t)}\le D_0\exp\left[\int_0^t(\|d\|^6_{L_3(S_2)}+\|\nabla v\|^2_{L_3(\Omega)})dt\right]\Lambda^2$$

and using Lemma 3.1 yields (3.34). □

Lemma 3.9. *Let the Conditions 1 and 2 hold. Then for a weak solution v to (1.1) and $h=v_{,x_3}$ - a solution to (3.29) there exists such constant $\mathcal{A}$ that*

$$\begin{aligned}&\mathbf{H}(T)\equiv\|h\|_{W_2^{2,1}(\Omega^T)}\le\mathcal{A}\\ &\text{and}\\ &\|v\|_{W_2^{2,1}(\Omega^T)}\le cD_0^2(\mathcal{A}^2+\Lambda^2)+c(\mathcal{A}+\mathcal{A}^2)+c\Lambda_2^2+c\Lambda_3^2+c\Lambda_4\end{aligned}\tag{3.35}$$

Proof. We note that

$$\|\nabla v\|_{L_2(0,T;L_3(\Omega))}\le c\|v\|_{W_{5/3}^{2,1}(\Omega^T)}\le cD_0(H_1+\Lambda)+c(H_1+H_1^2)+c\Lambda_2+c\Lambda_3$$

where we used (3.32). Applying also the inequality (3.34) from Lemma 3.8 we have

$$\|h\|^2_{L_2(\Omega^T)}\le D_0\exp\left[\int_0^t(\|d\|^6_{L_3(S_2)}dt+cD_0(H_1+\Lambda)+c(H_1+H_1^2)+c\Lambda_2+c\Lambda_3\right]\Lambda^2$$

Since

$$H_1(T)\le c\mathbf{H}(T)$$

we obtain, by formulas above

$$\begin{aligned}\mathbf{H}^2(T)\le cD_0^2\Lambda^2\left[(\mathbf{H}(T)+\Lambda)+c(\mathbf{H}(T)+\mathbf{H}(T)^2)+c\Lambda_2+c\Lambda_3\right]&\\ \cdot\exp\left[\int_0^t(\|d\|^6_{L_3(S_2)}dt+cD_0(\mathbf{H}(T)+\Lambda)+c(\mathbf{H}(T)+\mathbf{H}(T)^2)+c\Lambda_2+c\Lambda_3\right]&\\ +\|g\|^2_{L_2(\Omega^T)}+\|h(0)\|^2_{H^1(\Omega)}&\end{aligned}$$

To obtain the result, we need

$$\begin{aligned}
&cD_0^2\Lambda^2\left[(\mathcal{A}+\Lambda)+c(\mathcal{A}+\mathcal{A}^2)+c\Lambda_2+c\Lambda_3\right]\\
&\exp\left[\int_0^t(\|d\|^6_{L_3(S_2)}dt+cD_0(\mathcal{A}+\Lambda)+c(\mathcal{A}+\mathcal{A}^2)+c\Lambda_2+c\Lambda_3\right]\\
&+\|g\|^2_{L_2(\Omega^T)}+\|h(0)\|^2_{H^1(\Omega)}\le\mathcal{A}^2
\end{aligned}\tag{3.36}$$

which can be satisfied for sufficiently small $\|g\|_{L_2(\Omega^T)}, \|h(0)\|_{H^1(\Omega)}$ and

$$\Lambda=D_1+D_2+\|f_3\|^2_{L_2(0,t;L_{4/3}(S_2))}+\|g\|^2_{L_2(0,t;L_{6/5}(\Omega))}+\|h(0)\|^2_{L_2(\Omega)},$$

where

$$\begin{aligned}
D_1&=\|d_t\|^2_{L_2(0,t;H^1(S_2))}+\|d_{,x'}\|^2_{L_2(0,t;H^1(S_2))},\\
D_2&=\|d_{,x'}\|^2_{L_\infty(0,t;H^1(S_2))}.
\end{aligned}$$

To obtain the bound for v in terms of $\mathcal{A}$ we use (3.33) and $(3.35)_1$. This concludes the proof. □

4. Existence

Lemma 4.1. *Assume that Conditions 1 and 2 are satisfied. Then the solution* (v,p) *to (1.1) satisfying (1.8) exists on* $(0,T)$ *for some* $T>0$.

Proof. To prove the existence of solutions to problem (1.1) we will use the Leray-Schauder theorem. To this end, we construct the mappings

$$\begin{aligned}
&v_t-\operatorname{div}\mathbb{T}(v,p)=-\lambda\tilde v\cdot\nabla\tilde v+f && \text{in } \Omega^T=\Omega\times(0,T),\\
&\operatorname{div}v=0 && \text{in } \Omega^T,\\
&v\cdot\bar n=0 && \text{on } S_1^T,\\
&\nu\bar n\cdot\mathbb{D}(v)\cdot\bar\tau_\alpha+\gamma v\cdot\bar\tau_\alpha=0,\ \alpha=1,2, && \text{on } S_1^T,\\
&v\cdot\bar n=d && \text{on } S_2^T,\\
&\bar n\cdot\mathbb{D}(v)\cdot\bar\tau_\alpha=0,\quad \alpha=1,2, && \text{on } S_2^T,\\
&v\big|_{t=0}=v(0) && \text{in } \Omega,
\end{aligned}\tag{4.1}$$

and

$$\begin{aligned}
h_{,t}-\operatorname{div}\mathbb{T}(h,q)&=-\lambda(\tilde v\cdot\nabla\tilde h+\tilde h\cdot\nabla\tilde v)+g && \text{in } \Omega^T,\\
\operatorname{div}h&=0 && \text{in } \Omega^T,\\
n\cdot\bar h=0,\quad \bar n\cdot\mathbb{D}(h)\cdot\bar\tau_\alpha&+\gamma h\cdot\bar\tau_\alpha=0,\quad\alpha=1,2 && \text{on } S_1^T,\\
h_i=-d_{x_i},\quad i=1,2,\quad h_{3,x_3}&=\Delta'd && \text{on } S_2^T,\\
h\big|_{t=0}&=h(0) && \text{in } \Omega.
\end{aligned}\tag{4.2}$$

where $g=f_{,x_3}, \Delta'=\partial^2_{x_1}+\partial^2_{x_2}$, $\lambda\in[0,1]$ and $\tilde v$, $\tilde h$ are treated as given functions. We assume that $\tilde h=\tilde v_{,x_3}$, thus differentiating (4.1) with respect to x_3 and subtracting from (4.2) we obtain that

$$h=v_{,x_3}.$$

Problems (4.1), (4.2) define the mappings

$$\Phi_1 : \ (\tilde v,\lambda)\to(v,p),$$
$$\Phi_2 : \ (\tilde v,\tilde h,\lambda)\to(h,q).$$

We set $\Phi=(\Phi_1,\Phi_2)$. In previous Section we have shown a-priori estimate for a fixed point of Φ for $\lambda=1$. On the other hand, for $\lambda=0$ we have a unique existence of solutions to problems (4.1) and (4.2).

Let us introduce the space

$$\mathcal{M}(\Omega^T)=L_{2r}(0,(T;W^2_{\frac{6\eta}{3+\eta}}(\Omega)),\quad \eta\ge 2,\quad r\ge 2.$$

We shall find restrictions on r,η such that

$$\Phi:\ \mathcal{M}(\Omega^{(T})\times\mathcal{M}(\Omega^T)\to\mathcal{M}(\Omega^T)\times\mathcal{M}(\Omega^T)$$

is a compact mapping.
Assume that $\tilde v\in L_{2r}(0,T;W^1_{\frac{6\eta}{3+\eta}}(\Omega))$. Then

$$\begin{aligned}\|\tilde v\cdot\nabla\tilde v\|_{L_r(0,T;L_\eta(\Omega))}&=\bigg(\int_0^T dt\|\tilde v\cdot\nabla\tilde v\|^r_{L_\eta(\Omega)}\bigg)^{1/r}\\ &\le\bigg(\int_0^T dt\|\tilde v\|^r_{L_{\frac{6\eta}{3-\eta}}(\Omega)}\|\nabla\tilde v\|^r_{L_{\frac{6\eta}{3+\eta}}(\Omega)}\bigg)^{1/r}\\ &\le c\bigg(\int_0^T dt\|\tilde v\|^{2r}_{W^1_{\frac{6\eta}{3+\eta}}(\Omega)}\bigg)^{1/r}\le c\|\tilde v\|^2_{L_{2r}(0,T;W^1_{\frac{6\eta}{3+\eta}}(\Omega))}\end{aligned}\tag{4.3}$$

In the same way we obtain

$$\begin{aligned}&\|\tilde v\cdot\nabla\tilde h\|_{L_r(0,T;L_\eta(\Omega))}+\|\tilde h\cdot\nabla\tilde v\|_{L_r(0,T;L_\eta(\Omega))}\\ &\le c\|\tilde v\|_{L_{2r}(0,T;W^1_{\frac{6\eta}{3+\eta}}(\Omega))}\|\tilde h\|_{L_{2r}(0,T;W^1_{\frac{6\eta}{3+\eta}}(\Omega))}.\end{aligned}\tag{4.4}$$

In view of (4.3) and (4.4) can be shown that solutions to problems (4.1) and (4.2) belong to $W^{2,1}_{\eta,r}(\Omega^T)$ (see [S1]).
We are going to use the following imbeddings

$$W^{2,1}_2(\Omega^T)\supset W^{2,1}_{\eta,r}(\Omega^T)\tag{4.5}$$

and

$$W^{2,1}_2(\Omega^T)\subset L_{2r}(0,T;W^1_{\frac{6\eta}{3+\eta}}(\Omega))\equiv\mathcal{M}(\Omega^T),\tag{4.6}$$

where (4.5) holds for $\eta\ge 2$, $r\ge 2$ and (4.6) is compact for r,η satisfying the inequality

$$\frac52-\frac{3}{\frac{6\eta}{3+\eta}}-\frac{2}{2r}<1$$

which takes the form

$$1<\frac{3}{2\eta}+\frac1r.\tag{4.7}$$

Setting $r=\eta=2$ we obtain that $\tilde v,\tilde h\in L_4(0,T;W^1_{12/5}(\Omega))$ and then condition (4.7) takes the form

$$1<\frac34+\frac12\quad\text{so}\quad\frac12<\frac34.\tag{4.8}$$

Hence, we have compactness of mappings Φ_1 and Φ_2.
To show continuity of mappings Φ_1 and Φ_2 we consider

$$\begin{aligned}
&v_{st} - \operatorname{div}\mathbb{T}(v_s,p_s) = -\lambda\tilde{v}_s\cdot\nabla\tilde{v}_s + f && \text{in } \ \Omega^T,\\
&\operatorname{div} v_s = 0 && \text{in } \ \Omega^T,\\
&\bar{n}\cdot v_s = 0, \ \ \nu\bar{n}\cdot\mathbb{D}(v_s)\cdot\bar{\tau}_\alpha + \gamma v_s\cdot\bar{\tau}_\alpha = 0 && \text{on } \ S_1^T,\\
&v_s\cdot\bar{n} = d && \text{on } S_2^T,\\
&\bar{n}\cdot\mathbb{D}(v_s)\cdot\bar{\tau}_\alpha = 0, \quad \alpha = 1,2, && \text{on } \ S_2^T,\\
&v_s|_{t=0} = v(0) && \text{in } \ \Omega
\end{aligned}$$

and

$$\begin{aligned}
&h_{st} - \operatorname{div}\mathbb{T}(h_s,q_s) = -\lambda(\tilde{h}_s\cdot\nabla\tilde{v}_s + \tilde{v}_s\cdot\nabla\tilde{h}_s) + g && \text{in } \ \Omega^T\\
&\operatorname{div} h_s = 0 && \text{in } \ \Omega^T\\
&\bar{n}\cdot h_s = 0, \ \ \nu\bar{n}\cdot\mathbb{D}(h_s)\cdot\bar{\tau}_\alpha + \gamma h_s\cdot\bar{\tau}_\alpha = 0, \ \ \alpha = 1,2 && \text{on } \ S_1^T,\\
&h_{si} = -d_{x_i}, \ \ i = 1,2, \ \ h_{s3,x_3} = \Delta' d && \text{on } S_2^T,\\
&h_s|_{t=0} = h(0) && \text{in } \ \Omega,
\end{aligned}$$

where $s = 1,2$.
Let

$$V = v_1 - v_2, \quad H = h_1 - h_2, \quad P = p_1 - p_2, \quad Q = q_1 - q_2$$

Then V and H are solutions to the problems

$$\begin{aligned}
&V_t - \operatorname{div}\mathbb{T}(V,P) = -\lambda(\tilde{V}\cdot\nabla\tilde{v}_1 + \tilde{v}_2\cdot\nabla\tilde{V})\\
&\operatorname{div} V = 0\\
&V\cdot\bar{n}|_{S_1} = 0, \ \ \nu\bar{n}\cdot\mathbb{D}(V)\cdot\bar{\tau}_\alpha + \gamma V\cdot\bar{\tau}_\alpha|_{S_1} = 0, \ \ \alpha = 1,2,\\
&V\cdot\bar{n}|_{S_2} = d \ \ ,\bar{n}\cdot\mathbb{D}(V)\cdot\bar{\tau}_\alpha|_{S_2} = 0, \ \ \alpha = 1,2,\\
&V|_{t=0} = 0,
\end{aligned}\tag{4.9}$$

$$\begin{aligned}
&H_t - \operatorname{div}\mathbb{T}(H,Q) = -\lambda(\tilde{H}\cdot\nabla\tilde{v}_1 + \tilde{h}_2\cdot\nabla\tilde{V} + \tilde{V}\cdot\nabla\tilde{h}_1 + \tilde{v}_2\cdot\nabla\tilde{H})\\
&\operatorname{div} H = 0\\
&H\cdot\bar{n}|_{S_1} = 0, \ \ \nu\bar{n}\cdot\mathbb{D}(H)\cdot\bar{\tau}_\alpha + \gamma H\cdot\bar{\tau}_\alpha|_{S_1} = 0, \ \ \alpha = 1,2,\\
&H_i|_{S_2} = -d_{x_i}, \ \ i = 1,2, \ \ H_{3,x_3}|_{S_2} = \Delta' d,\\
&H|_{t=0} = 0.
\end{aligned}\tag{4.10}$$

Assume that $\lambda \neq 0$. In Lemma 3.9, we proved the existence of such a constant $\mathcal{A}$ that for sufficiently chosen data,

$$\|h\|_{W_2^{2,1}(\Omega^T)} \le \mathcal{A} \quad \text{and} \quad \|v\|_{W_2^{2,1}(\Omega^T)} \le \varphi(\mathcal{A}).$$

Then for solutions of (4.9) we have

$$\begin{aligned}
\|V\|_{\mathcal{M}(\Omega^T)} &= \|V\|_{L_{2r}(0,T;W^1_{\frac{6\eta}{3+\eta}}(\Omega))} \le c\|V\|_{W_2^{2,1}(\Omega^T)} \le c\|V\|_{W^{2,1}_{\eta,r}(\Omega^T)}\\
&\le c\sum_{s=1}^{2}\|\tilde{v}_s\|_{L_{2r}(0;T;W^1_{\frac{6\eta}{3+\eta}}(\Omega))}\cdot\|\tilde{V}\|_{L_{2r}(0;T;W^1_{\frac{6\eta}{3+\eta}}(\Omega))} \le c(\mathcal{A})\|\tilde{V}\|_{\mathcal{M}(\Omega^T)},
\end{aligned}\tag{4.11}$$

where $r \ge 2$, $\eta \ge 2$ and satisfy either (4.7) or (4.8).

Similarly we can prove that

$$\|H\|_{\mathcal{M}(\Omega^T)} \le c(\mathcal{A})(\|\tilde{V}\|_{\mathcal{M}(\Omega^T)} + \|\tilde{H}\|_{\mathcal{M}(\Omega^T)}) \tag{4.12}$$

Inequalities (4.11) and (4.12) imply continuity of mapping Φ. Continuity with respect to λ is obvious. Hence by the Leray-Schauder fixed point theorem we have the existence of solutions to problem (1.1) such that $v \in W_2^{2,1}(\Omega^T)$, $\nabla p \in L_2(\Omega^T)$ and estimates (1.8) hold. □

Lemma 4.1 and estimates (3.35) from Lemma 3.9 imply Theorem 1.

References

[BIN] O. V. Besov, V. P. Il'in and S. M. Nikol'skii, *Integral representations of functions and imbedding theorems.*, Vol. I. Translated from the Russian. Scripta Series in Mathematics, New York-Toronto, Ont.-London, 1978. viii+345 pp.

[L] O. A. Ladyzhenskaya, *Mathematical Problems of Viscous Incompressible Fluid*, Nauka, Moscow 1970 (in Russian).

[M1] P. B. Mucha, *On the inviscid limit of the Navier-Stokes equations for flows with large flux*, Nonlinearity 16, No 5, 1715-1732 (2003).

[M2] P. B. Mucha, *On cylindrical symmetric flows through pipe-like domains*, Journal of Differential Equations 201, No 2, 304-323 (2004).

[RZ1] J. Rencławowicz and W.M. Zajączkowski, *Existence of global weak solutions for Navier-Stokes equations with large flux.*, ArXiv no 1009.5699.

[RZ2] J. Rencławowicz and W.M. Zajączkowski, *Large time regular solutions to the Navier-Stokes equations in cylindrical domains*, Topol. Methods Nonlinear Anal. 32 (2008),69-87.

[S1] V.A. Solonnikov, *Solvability of the Stokes equations in S. L. Sobolev spaces with a mixed norm*, Zap. Nauchn. Sem. POMI 288 (2002), 204–231.

[Z1] W.M. Zajączkowski, *Global regular nonstationary flow for the Navier-Stokes equations in a cylindrical pipe*, TMNA 26(2005), 221-286.

[Z2] W. M. Zajączkowski, *Global special regular solutions to the Navier-Stokes equations in a cylindrical domain without the axis of symmetry*, Top. Meth. Nonlinear Anal. 24 (2004), 69-105.

[Z3] W. M. Zajączkowski, *On global regular solutions to the Navier-Stokes equations in cylindrical domains*, to appear in TMNA.

[Z4] W. M. Zajączkowski, *Large time existence of solutions to the Navier-Stokes equations in axially symmetric domains with inflow and outflow*, Functiones et Approximatio 40.2 (2009), 209-250.

Joanna Rencławowicz: Institute of Mathematics, Polish Academy of Sciences, Śniadeckich 8, 00-956 Warsaw, Poland, e-mail: jr@impan.gov.pl

Wojciech M. Zajączkowski: Institute of Mathematics, Polish Academy of Sciences, Śniadeckich 8, 00-956 Warsaw, Poland, e-mail: wz@impan.gov.pl and Institute of Mathematics and Cryptology, Military University of Technology, Kaliskiego 2, 00-908 Warsaw, Poland